

Weekly respiratory pathogens report

Week 19 of 2022

Highlights

- In 2022 to date, 106 sporadic influenza cases have been detected, with increasing number of cases reported in the past 3 weeks. Majority of cases were reported from, Kwa-Zulu Natal (n=41), followed by Mpumalanga (n=31), Gauteng (n=28), North West (n=4) and Western Cape (n=2) sentinel surveillance sites.
- The 2022 respiratory syncytial virus (RSV) season which started in week7 (week starting 14 February 2022) when RSV detection rate among children under five years of age in pneumonia surveillance rose above the seasonal threshold, continues. In Week 19, RSV activity among children aged < 5 years was on moderate threshold.
- In 2022 to date, a total of 375 COVID-19 cases were detected from all surveillance programmes. In Week 19, an increase in detection rate of COVID-19 cases has been noted in both influenza-like illness (ILI) programme and pneumonia surveillance. Of the 187 hospitalised COVID-19 cases reported with available data on outcome, 13 (7%) died.
- Of the 284/375 (76%) with variant data from all surveillance programmes, Omicron was the dominant variant 50% (144/284) of which 58% (76/144) was Omicron (21K/BA.1) sub-lineage, 45% (65/144) was Omicron (21L/BA.2) sub-lineage, 1% (2/144) was Omicron (21M/BA.3) sub-lineage and <1% (1/144) was Omicron (22A/BA.4) sub-lineage. Alpha (1/284) and Delta (1/284) variants contributed <1% each and for 49% (138/284) variant was not assigned.

Programme Descriptions

Programme	Influenza-like illness (ILI)	Viral Watch	National syndromic surveillance for pneumonia
Start year	2012	1984	2009
Provinces*	KZ NW WC MP	EC FS GP LP MP NC NW WC	EC GP KZ MP NW WC
Type of site	Primary health care clinics	General practitioners	Public hospitals
Case definition	<p>ILI: An acute respiratory illness with a temperature ($\geq 38^{\circ}\text{C}$) and cough, & onset ≤ 10 days</p> <p>Suspected pertussis Any person with an acute cough illness lasting ≥ 14 days (or cough illness of any duration for children < 1 year), without a more likely diagnosis AND one or more of the following signs or symptoms:</p> <ul style="list-style-type: none"> • paroxysms of coughing, • or inspiratory "whoop", • or post-tussive vomiting • or apnoea in children < 1 year; <p>OR</p> <p>Any person in whom a clinician suspects pertussis</p> <p>Suspected SARS-CoV-2 Any person presenting with an acute (≤ 14 days) respiratory tract infection or other clinical illness compatible with COVID-19[§]</p>	<p>ILI: An acute respiratory illness with a temperature ($\geq 38^{\circ}\text{C}$) and cough, & onset ≤ 10 days</p> <p>Suspected SARS-CoV-2 Any person presenting with an acute (≤ 14 days) respiratory tract infection or other clinical illness compatible with COVID-19[§]</p>	<p>SRI: Acute (symptom onset ≤ 10 days) or chronic (symptom onset > 10) lower respiratory tract infection</p> <p>Suspected pertussis Any person with an acute cough illness lasting ≥ 14 days (or cough illness of any duration for children < 1 year), without a more likely diagnosis AND one or more of the following signs or symptoms:</p> <ul style="list-style-type: none"> • paroxysms of coughing, • or inspiratory "whoop", • or post-tussive vomiting • or apnoea in children < 1 year; <p>OR</p> <p>Any person in whom a clinician suspects pertussis.</p> <p>Suspected SARS-CoV-2 Any person admitted with a physician-diagnosis of suspected COVID-19 and not meeting SRI case definition.</p>
Specimens collected	Oropharyngeal & nasopharyngeal swabs	Throat and/or nasal swabs or Nasopharyngeal swabs	Oropharyngeal & nasopharyngeal swabs
Main pathogens tested**	INF RSV BP SARS-CoV-2	INF RSV BP SARS-CoV-2	INF RSV BP SARS-CoV-2
Testing Methods	<p>INF and RSV - Fast-Track Diagnostics multiplex real-time reverse transcription polymerase chain reaction (until 31 March 2021)</p> <p>B. pertussis Multiplex real-time PCR (Tatti <i>et al.</i>, <i>J Clin Microbiol</i> 2011) and culture (if PCR cycle threshold ≤ 25)</p> <p>SARS-CoV-2 1 April 2020 – 31 March 2021: Roche E gene real-time PCR assay (Corman <i>et al.</i>, <i>Euro Surv</i> 2020) 1 April 2021 to date: Allplex™ SARS-CoV-2/FluA/FluB/RSV PCR kit</p> <p>- positivity assigned if PCR cycle threshold is < 40 for ≥ 1 gene targets (N, S, OR RdRp)</p>	<p>INF and RSV - Fast-Track Diagnostics multiplex real-time reverse transcription polymerase chain reaction (until 31 March 2021)</p> <p>B. pertussis Multiplex real-time PCR (Tatti <i>et al.</i>, <i>J Clin Microbiol</i> 2011) and culture (if PCR cycle threshold ≤ 25)</p> <p>SARS-CoV-2 1 April 2020 – 31 March 2021: Roche E gene real-time PCR assay (Corman <i>et al.</i>, <i>Euro Surv</i> 2020) 1 April 2021 to date: Allplex™ SARS-CoV-2/FluA/FluB/RSV PCR kit</p> <p>- positivity assigned if PCR cycle threshold is < 40 for ≥ 1 gene targets (N, S, OR RdRp)</p>	<p>INF and RSV - Fast Track Diagnostics multiplex real-time reverse transcription polymerase chain reaction (until 31 March 2021)</p> <p>B. pertussis Multiplex real-time PCR (Tatti <i>et al.</i>, <i>J Clin Microbiol</i> 2011) and culture (if PCR cycle threshold ≤ 25)</p> <p>SARS-CoV-2 1 April 2020 – 31 March 2021: Roche E gene real-time PCR assay (Corman <i>et al.</i>, <i>Euro Surv</i> 2020) 1 April 2021 to date: Allplex™ SARS-CoV-2/FluA/FluB/RSV PCR kit</p> <p>- positivity assigned if PCR cycle threshold is < 40 for ≥ 1 gene targets (N, S, OR RdRp)</p>

Epidemic Threshold

Thresholds are calculated using the Moving Epidemic Method (MEM), a sequential analysis using the R Language, available from: <http://CRAN.R-project.org/web/package=mem> designed to calculate the duration, start and end of the annual influenza epidemic. MEM uses the 40th, 90th and 97.5th percentiles established from available years of historical data to calculate thresholds of activity. Thresholds of activity for influenza and RSV are defined as follows: Below seasonal threshold, Low activity, Moderate activity, High activity, Very high activity. For influenza, thresholds from outpatient influenza like illness (ILI in primary health care clinics) are used as an indicator of disease transmission in the community and thresholds from pneumonia surveillance are used as an indicator of impact of disease. For RSV, thresholds from pneumonia surveillance, using data from children aged < 5 years are used to define the start and end of the season.

* EC: Eastern Cape; FS: Free State; GP: Gauteng; KZ: KwaZulu-Natal; LP: Limpopo; MP: Mpumalanga; NC: Northern Cape; NW: North West; WC: Western Cape

**INF: influenza virus; RSV: respiratory syncytial virus; BP: *Bordetella pertussis*; SARS-CoV-2: severe acute respiratory syndrome coronavirus 2

[§]Symptoms include ANY of the following respiratory symptoms: cough, sore throat, shortness of breath, anosmia (loss of sense of smell) or dysgeusia (alteration of the sense of taste), with or without other symptoms (which may include fever, weakness, myalgia, or diarrhoea). Testing for SARS-CoV-2 was initiated in all three surveillance programmes in week 10 of 2020 (week starting 2 March 2020).




Figure 1. Number of influenza positive cases* by influenza subtype and lineage and detection rate*** by week, Influenza-like illness (ILI) surveillance in primary health care clinics, 03/01/2022 – 15/05/2022**

*Specimens from patients with influenza-like illnesses at 5 sentinel sites in 4 provinces

***Only reported for weeks with >10 specimens submitted

Inconclusive: insufficient viral load in sample and unable to characterise further

**Influenza was detected in five (16%) of 31 specimens from patients who met suspected SARS-CoV-2 case definition but did not meet Influenza-like illness (ILI) case definition. Of which four (80%) were influenza A(H1N1)pdm09, and one (20%) was influenza B(Victoria). These are not included in the epidemiological curve.

Table 1. Number of laboratory confirmed influenza cases by subtype and lineage and total number of samples tested by clinic and province, Influenza-like illness (ILI) surveillance in primary health care clinics, 03/01/2022 – 15/05/2022**

Clinic (Province)	A(H1N1)pdm09	A(H3N2)	A subtype inconclusive	A subtype pending results ^β	B/Victoria	B/Yamagata	B lineage inconclusive	B lineage pending results ^β	Total samples
Agincourt (MP)	13	0	0	0	0	0	0	0	99
Eastridge (WC)	0	0	0	0	0	0	0	0	92
Edendale	14	0	0	3	0	0	0	0	168
Gateway (KZ)									
Jouberton (NW)	1	0	0	1	0	0	0	0	158
Mitchell's Plain (WC)	0	0	1	0	0	0	0	0	52
Total:	28	0	1	4	0	0	0	0	569

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape; MP: Mpumalanga

Inconclusive: insufficient viral load in sample and unable to characterise further

^βinfluenza A subtype or B lineage results are pending

**Influenza was detected in five (16%) of 31 specimens from patients who met suspected SARS-CoV-2 case definition but did not meet Influenza-like illness (ILI) case definition. Of which four (80%) were influenza A(H1N1)pdm09, and one (20%) was influenza B(Victoria). These are not included in the epidemiological curve.




Figure 2. Influenza percentage detections and epidemic thresholds* among cases of all ages, Influenza-like illness (ILI) surveillance in primary health care clinics, 03/01/2022 – 15/05/2022

*Thresholds based on 2012-2019 data




Figure 3. Number of patients testing positive for respiratory syncytial virus* by province and detection rate by week, Influenza-like illness (ILI) surveillance in primary health care clinics, 03/01/2022 – 15/05/2022

*RSV was not detected from 31 specimens of patients who met suspected SARS-CoV-2 case definition but did not meet influenza-like illness (ILI) case definition.




Figure 4. Number of patients testing positive for respiratory syncytial virus* by subgroup and detection rate by week, Influenza-like illness (ILI) surveillance in primary health care clinics, 03/01/2022 – 15/05/2022

Inconclusive: insufficient viral load in sample and unable to characterise further

RSV AB: Both RSV A and B subgroup identified.

*RSV was not detected from 31 specimens of patients who met suspected SARS-CoV-2 case definition but did not meet influenza-like illness (ILI) case definition. These are not included in the epidemiological curve.

Table 2. Number of patients testing positive for respiratory syncytial virus (RSV) by subgroups identified and total number of samples tested by clinic and province, Influenza-like illness (ILI) surveillance in primary health care clinics, 03/01/2022 – 15/05/2022**

Clinic (Province)	RSVA	RSVB	RSVAB	RSV subgroup inconclusive	RSV subgroup pending*	Total samples
Agincourt (MP)	16	1	0	1	0	99
Eastridge (WC)	0	3	0	0	2	92
Edendale Gateway (KZ)	26	0	0	3	0	168
Jouberton (NW)	1	3	0	0	1	158
Mitchell's Plain (WC)	0	1	0	0	0	52
Total	43	8	0	4	3	569

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape; MP: Mpumalanga

Inconclusive: insufficient viral load in sample and unable to characterise further

RSV AB: Both RSV A and B subgroup identified

*RSV results for subgroups are pending

**RSV was not detected from 31 specimens of patients who met suspected SARS-CoV-2 case definition but did not meet influenza-like illness (ILI) case definition. These are not included in the table.




Figure 5. Number of patients testing positive for SARS-CoV-2* by province and detection rate by week, Influenza-like illness (ILI) surveillance in primary health care clinics, 03/01/2022 – 15/05/2022

*Specimens from patients with influenza-like illnesses at 5 sentinel sites in 4 provinces

*SARS-CoV-2 was detected in 6 of 31 (13%) specimens from patients who met suspected SARS-CoV-2 case definition but did not meet influenza-like illness (ILI) case definition. These are not included in the epidemiological curve.

Table 3. Number of patients positive for SARS-CoV-2* identified and total number of samples tested by clinic and province, Influenza-like illness (ILI) surveillance primary health care clinics, 03/01/2022 – 15/05/2022

Clinic (Province)	SARS-CoV-2 positive	Total samples tested
Agincourt (MP)	25	99
Eastridge (WC)	5	92
Edendale Gateway (KZ)	11	168
Jouberton (NW)	34	158
Mitchell's Plain (WC)	18	52
Total:	93	569

KZ: KwaZulu-Natal; NW: North West; WCP: Western Cape; MP: Mpumalanga

*SARS-CoV-2 was detected in 6 of 31 (13%) specimens from patients who met suspected SARS-CoV-2 case definition but did not meet influenza-like illness (ILI) case definition. These are not included in the table.




Figure 6. Number and detection rate of laboratory confirmed SARS-CoV-2* cases by variant type (variant PCR/sequencing) and week, Influenza-like illness (ILI) surveillance in primary health care clinics, 03/01/2022 – 15/05/2022

*Specimens are from patients with influenza-like illness at 5 sentinel sites in 4 provinces who met suspected SARS-CoV-2 case definition or met ILI case definition

Unable to assign: no lineage assigned due to poor- sequence quality OR low viral load (ct=>35) OR variant PCR could not assign variant and no sequencing result

Pending: outstanding variant results

Table 4. Number of cases tested and number positive for SARS-CoV-2* by variant (variant PCR and/or sequencing) identified and total number of samples tested by clinic and province, Influenza-like illness (ILI) surveillance primary health care clinics, 03/01/2022 – 15/05/2022

Clinic (Province)	Non-Alpha/Beta/Delta	Alpha (20I)	Beta (20H)	Delta (21A, 21I, 21J)	C.1.2 (20D)	Omicron (21K/BA.1)	Omicron (21L/BA.2)	Omicron (21M/BA.3)	Omicron (22A/BA.4)	Unable to assign	Pending	Total SARS-CoV-2 positive	Total samples tested
Agincourt (MP)	0	0	0	0	0	4	3	0	0	20	0	27	107
Eastridge (WC)	0	0	0	0	0	2	0	0	0	1	2	5	92
Edendale	0	0	0	0	0	2	1	0	0	4	6	13	184
Gateway (KZ)	0	0	0	0	0	1	5	0	0	21	9	36	165
Jouberton (NW)	0	0	0	0	0	2	0	0	0	4	12	18	52
Mitchell's Plain (WC)	0	0	0	0	0	11	9	0	0	50	29	99	600

KZ: KwaZulu-Natal; NW: North West; WCP: Western Cape; MP: Mpumalanga

*Specimens are from patients with influenza-like illness at 5 sentinel sites in 4 provinces who met suspected SARS-CoV-2 case definition or met ILI case definition

Unable to assign: no lineage assigned due to poor- sequence quality OR low viral load (ct=>35) OR variant PCR could not assign variant and no sequencing result

Pending: outstanding variant results




Figure 7. Number of positive patients* by influenza subtype and lineage and detection rate** by week, ILI surveillance - Viral Watch, 03/01/2022 – 15/05/2022

*Specimens from patients with Influenza-like illnesses at 90 sentinel sites in 8 provinces

** Only reported for weeks with >10 specimens submitted.

Inconclusive: insufficient viral load in sample and unable to characterise further

Table 5. Number of laboratory confirmed influenza cases by influenza subtype and lineage and total number of samples tested by province, ILI surveillance - Viral Watch, 03/01/2022 – 15/05/2022

Province	A(H1N1) pdm09	A(H3N2)	A subtype inconclusiv e	A subtype pending results*	B/Victor ia	B/Yamag ata	B lineage inconclus ive	B lineage pending results*	Total samples
Eastern Cape	0	0	0	0	0	0	0	0	2
Free State	0	0	0	0	0	0	0	0	0
Gauteng	3	1	0	9	0	0	0	1	92
Limpopo	0	0	0	0	0	0	0	0	0
Mpumalanga	0	0	0	0	0	0	0	0	2
North West	0	0	0	0	0	0	0	0	2
Northern Cape	0	0	0	0	0	0	0	0	0
Western Cape	0	0	0	0	0	0	0	0	32
Total:	3	1	0	9	0	0	0	1	130

Inconclusive: insufficient viral load in sample and unable to characterise further

*Influenza A subtype or B lineage results are pending




Figure 8. Number of RSV positive cases testing positive for respiratory syncytial virus (RSV)* by subgroup and detection rate** by week, ILI surveillance - Viral Watch, 03/01/2022 – 15/05/2022

*Specimens from patients with Influenza-like illnesses at 92 sentinel sites in 8 provinces

** Only reported for weeks with >10 specimens submitted.

Table 6. Number of RSV positive cases identified and total number of samples tested by province, ILI surveillance - Viral Watch, 03/01/2022 – 15/05/2022

Province	RSV A	RSV B	RSV AB	RSV subgroup inconclusive	RSV subgroup pending results*	Total samples tested
Eastern Cape	0	0	0	0	0	2
Free State	0	0	0	0	0	0
Gauteng	2	5	0	0	1	92
Limpopo	0	0	0	0	0	0
Mpumalanga	0	0	0	0	0	2
North West	0	0	0	0	0	2
Northern Cape	0	0	0	0	0	0
Western Cape	0	3	0	0	3	32
Total:	2	8	0	0	4	130

*RSV results for subgroups are pending

**Inconclusive: insufficient viral load in sample and unable to characterise further




Figure 9. Number of patients testing positive for SARS-CoV-2*, by site and detection rate by week, ILI surveillance - Viral Watch, 03/01/2022 – 15/05/2022**

*Specimens from patients with Influenza-like illnesses at 92 sentinel sites in 8 provinces

** Only reported for weeks with >10 specimens submitted.

Table 7. Number of SARS-CoV-2 positive cases identified and total number tested by province, ILI surveillance - Viral Watch, 03/01/2022 – 15/05/2022

Province	SARS-CoV-2 positive	Total samples tested
Eastern Cape	1	2
Free State	0	0
Gauteng	41	92
Limpopo	0	0
Mpumalanga	2	2
North West	0	2
Northern Cape	0	0
Western Cape	10	32
Total:	54	130




Figure 10. Number and detection rate of laboratory confirmed SARS-CoV-2* cases by variant type (variant PCR/sequencing) and week, ILI surveillance - Viral Watch, 03/01/2022 – 15/05/2022

*Specimens from patients with Influenza-like illnesses at 92 sentinel sites in 8 provinces

Unable to assign: no lineage assigned due to poor- sequence quality **OR** low viral load (ct=>35) **OR** variant PCR could not assign variant and no sequencing result

Pending: outstanding variant results

Table 8. Number of SARS-CoV-2* positive cases by variant (variant PCR and/or sequencing) identified and total number of samples tested by province, ILI surveillance - Viral Watch, 03/01/2022 – 15/05/2022

Clinic (Province)	Non-Alpha/Beta/Delta	Alpha (20I)	Beta (20H)	Delta (21A,21I, 21J)	C.1.2 (20D)	Omicron (21K/BA.1)	Omicron (21L/BA.2)	Omicron (22A/BA.4)	Unable to assign	Pending	Total SARS-CoV-2 positive	Total samples tested
Eastern Cape	0	0	0	0	0	0	1	0	0	0	1	2
Free State	0	0	0	0	0	0	0	0	0	0	0	0
Gauteng	0	0	0	0	0	2	8	1	10	20	41	92
Limpopo	0	0	0	0	0	0	0	0	0	0	0	0
Mpumalanga	0	0	0	0	0	0	0	0	0	2	2	2
North West	0	0	0	0	0	0	0	0	0	0	0	2
Northern Cape	0	0	0	0	0	0	0	0	0	0	0	0
Western Cape	0	0	0	0	0	0	6	0	2	2	10	32
Total:	0	0	0	0	0	2	15	1	12	24	54	130

*Specimens from patients with Influenza-like illnesses at 92 sentinel sites in 8 provinces

Unable to assign: no lineage assigned due to poor- sequence quality **OR** low viral load (ct=>35) **OR** variant PCR could not assign variant and no sequencing result

Pending: outstanding variant results




Figure 11. Number of positive influenza positive cases* by influenza subtype and lineage and detection rate*** by week, pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022**

Inconclusive: insufficient viral load in sample and unable to characterise further

*Specimens from patients hospitalised with pneumonia at 7 sentinel sites in 5 provinces

***Only reported for weeks with >10 specimens submitted

**Influenza was not detected in 11 specimens from patients who met suspected SARS-CoV-2 case definition but did not meet pneumonia (SRI) case definition. These are not included in the epidemiological curve.

Table 9. Number of laboratory confirmed influenza cases by subtype and lineage* and total number of samples tested by hospital, pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022

Hospital (Province)	A(H1N1)pdm09	A(H3N2)	A subtype inconclusive	A subtype pending results***	B/Victoria	B/Yamagata	B lineage inconclusive	B lineage pending results***	Total samples
Edendale (KZ)	22	0	0	0	0	0	0	0	428
Helen Joseph-Rahima Moosa (GP)	7	1	0	4	0	0	0	0	631
Klerksdorp-Tshepong (NW)	1	0	0	1	0	0	0	0	206
Livingstone (EC)	0	0	0	0	0	0	0	0	12
Mapulaneng-Matikwana (MP)	7	0	0	0	0	0	0	0	202
Red Cross (WC)	1	0	0	0	0	0	0	0	272
Mitchell's Plain (WC)	0	0	0	0	0	0	0	0	453
Tembisa (GP)	1	0	0	1	0	0	0	0	53
Tintswalo (MP)	7	1	0	0	0	0	0	0	127
Tygerberg (WC)	0	0	0	0	0	0	0	0	6
Total:	46	2	0	6	0	0	0	0	2390

EC: Eastern Cape (Livingstone started enrolling on the 3rd of May 2022); GP: Gauteng (Tembisa started enrolling on the 10th March 2022); KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape (Tygerberg started enrolling on the 20th April 2022)

Inconclusive: insufficient viral load in sample and unable to characterise further

***influenza A subtype or B lineage results are pending

*Influenza was not detected in 11 specimens from patients who met suspected SARS-CoV-2 case definition but did not meet pneumonia (SRI) case definition. These are not included in the table.




Figure 12. Influenza percentage detections and epidemic thresholds* among cases of all ages, pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022

*Thresholds based on 2010-2019 data




Figure 13. Number of patients (all ages) testing positive for respiratory syncytial virus* by province and detection rate by week, pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022

*RSV was not detected in 11 specimens from patients who met suspected SARS-CoV-2 case definition but did not meet pneumonia (SRI) case definition.




Figure 14. Number of patients (all ages) testing positive for respiratory syncytial virus* by subgroup and detection rate by week, pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022

Inconclusive: insufficient viral load in sample and unable to characterise further

RSV AB: Both RSV A and B subgroup identified

RSV subgroup pending: RSV results for subgroups are pending

*RSV was not detected in 11 specimens from patients who met suspected SARS-CoV-2 case definition but did not meet pneumonia (SRI) case definition. These are not included in the epidemiological curve.

Table 10. Number of patients (all ages) positive for respiratory syncytial virus subgroups by subgroups identified and total number of samples tested by hospital, pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022**

Hospital (Province)	RSVA	RSVB	RSVAB	RSV subgroup inconclusive	RSV subgroup pending*	Total samples
Edendale (KZ)	80	1	0	2	0	428
Helen Joseph-Rahima Moosa (GP)	31	132	3	1	3	631
Klerksdorp-Tshepong (NW)	16	21	0	0	9	206
Livingstone (EC)	0	0	0	1	2	12
Mapulaneng-Matikwana (MP)	8	5	0	0	0	202
Red Cross (WC)	2	32	0	0	4	272
Mitchell's Plain (WC)	24	95	0	2	17	453
Tembisa (GP)	0	1	0	0	0	53
Tintswalo (MP)	3	7	0	3	0	127
Tygerberg (WC)	0	0	0	0	0	6
Total:	164	294	3	9	35	2390

EC: Eastern Cape (Livingstone started enrolling on the 3rd of May 2022); GP: Gauteng (Tembisa started enrolling on the 10th March 2022); KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape (Tygerberg started enrolling on the 20th April 2022)

Inconclusive: insufficient viral load in sample and unable to characterise further

RSV AB: Both RSV A and B subgroup identified

*RSV results for subgroups are pending

**RSV was not detected in 11 specimens from patients who met suspected SARS-CoV-2 case definition but did not meet pneumonia (SRI) case definition. These are not included in the table.




Figure 15. RSV percentage detections and epidemic thresholds* among children aged < 5 years, pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022

*Thresholds based on 2010-2019 data




Figure 16. Number of patients testing positive for SARS-CoV-2* by province and detection rate by week, pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022

*Specimens from patients hospitalized with pneumonia at 6 sentinel sites in 5 provinces

*SARS-CoV-2 was detected in 6 of 11 (55%) specimens from patients who met suspected SARS-CoV-2 case definition but did not meet pneumonia (SRI) case definition. These are not included in the epidemiological curve.

Table 11. Number of patients positive for SARS-CoV-2* and total number of samples tested by hospital, pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022

Hospital (Province)	SARS-CoV-2 positive	Total samples tested
Edendale (KZ)	62	428
Helen Joseph-Rahima Moosa (GP)	37	630
Klerksdorp-Tshepong (NW)	27	206
Livingstone (EC)	2	12
Mapulaneng-Matikwana (MP)	25	202
Red Cross (WC)	27	272
Mitchell's Plain (WC)	20	453
Tembisa (GP)	4	53
Tintswalo (MP)	12	127
Tygerberg (WC)	0	6
Total:	216	2389

EC: Eastern Cape (Livingstone started enrolling on the 3rd of May 2022); GP: Gauteng (Tembisa started enrolling on the 10th March 2022); KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape (Tygerberg started enrolling on the 20th April 2022)

*SARS-CoV-2 was detected in 6 of 11 (55%) specimens from patients who met suspected SARS-CoV-2 case definition but did not meet pneumonia (SRI) case definition. These are not included in the table.




Figure 17. Number and detection rate of laboratory confirmed SARS-CoV-2 cases* by variant type (variant PCR/sequencing), pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022

*Specimens are from hospitalized patients at 7 sentinel sites in 5 provinces who met suspected SARS-CoV-2 case definition and met pneumonia (SRI) case definition

Unable to assign: no lineage assigned due to poor- sequence quality **OR** low viral load (ct=>35) **OR** variant PCR could not assign variant and no sequencing result

Pending: outstanding variant results

Table 12. Number of SARS-CoV-2 positive cases* by variant (variant PCR and/or sequencing) identified and total number of samples tested by hospital, pneumonia surveillance public hospitals, 03/01/2022 – 15/05/2022

Hospital (Province)	Non-Alpha/Beta/Delta	Alpha (20I)	Beta (20H)	Delta (21A, 21I, 21J)	C.1.2 (20D)	Omicron (21K/BA.1)	Omicron (21L/BA.2)	Omicron (21M/BA.3)	Unable to assign	Pending	Total SARS-CoV-2 positive	Total samples tested
Edendale (KZ)	0	0	0	1	0	24	13	1	17	10	66	436
Helen Joseph-Rahima Moosa (GP)	0	1	0	0	0	6	9	0	12	9	37	630
Klerksdorp-Tshepong (NW)	0	0	0	0	0	10	2	1	12	2	27	206
Livingstone (EC)	0	0	0	0	0	0	0	0	0	2	2	12
Mapulaneng-Matikwana (MP)	0	0	0	0	0	4	8	0	13	2	27	205
Red Cross (WC)	0	0	0	0	0	11	1	0	10	5	27	272
Mitchell's Plain (WC)	0	0	0	0	0	4	4	0	7	5	20	453
Tembisa (GP)	0	0	0	0	0	1	0	0	0	3	4	53
Tintswalo (MP)	0	0	0	0	0	3	4	0	5	0	12	127
Tygerberg (WC)	0	0	0	0	0	0	0	0	0	0	0	6
Total:	0	1	0	1	0	63	41	2	76	38	222	2400

EC: Eastern Cape (Livingstone started enrolling on the 3rd of May 2022); GP: Gauteng (Tembisa started enrolling on the 10th March 2022); KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape (Tygerberg started enrolling on the 20th April 2022)

*Specimens are from hospitalized patients at 7 sentinel sites in 5 provinces who met suspected SARS-CoV-2 case definition and met pneumonia (SRI) case definition

Unable to assign: no lineage assigned due to poor- sequence quality **OR** low viral load (ct=>35) **OR** variant PCR could not assign variant and no sequencing result

Pending: outstanding variant results

Summary of individuals with laboratory confirmed SARS-CoV-2

Table13: Characteristics of individuals with laboratory-confirmed SARS-CoV-2, enrolled in influenza-like illness (ILI) and pneumonia surveillance programmes, South Africa, 03 January 2022 – 15 May 2022

Characteristic	Influenza-like illness (ILI), public-sector, n=99 (%)	Pneumonia, n=222 (%)
Age group (years)		
0-9	16/99 (16)	55/222 (25)
10-19	10/99 (10)	6/222 (3)
20-39	23/99 (23)	53/222 (24)
40-59	27/99 (28)	56/222 (25)
60-79	7/99 (7)	36/222 (16)
≥80	1/99 (1)	7/222 (3)
Sex-female	52/99 (53)	111/222 (50)
Province*		
Eastern Cape	N/A	2/222 (1)
Gauteng	N/A	41/222 (18)
KwaZulu-Natal	13/99 (13)	66/222 (30)
Mpumalanga	27/99 (27)	39/222 (18)
North West	36/99 (37)	27/222 (12)
Western Cape	23/99 (23)	47/222 (21)
Race		
Black	62/99 (63)	169/222 (76)
Coloured	11/99 (11)	36/222 (16)
Asian/Indian	0/99 (0)	1/222 (0)
White	11/99 (11)	5/222 (2)
Other	15/99 (15)	11/222 (5)
Variant		
Non-Alpha/Beta/Delta	0/99 (0)	0/222 (0)
Alpha(20I)	0/99 (0)	1/222 (0)
Beta(20H)	0/99 (0)	0/222 (0)
Delta(21A, 21I, 21J)	0/99 (0)	1/222 (0)
C.1.2(20D)	0/99 (0)	0/222 (0)
Omicron (21K/BA.1)	11/99 (11)	63/222 (28)
Omicron (21L/BA.2)	9/99 (9)	41/222 (18)
Omicron (21M/BA.3)	0/99 (0)	2/222 (0)
Unable to assign ⁵⁵	50/99 (51)	76/222 (34)
Pending results ⁵	29/99 (30)	38/222 (17)
Presentation		
Fever	62/83 (75)	87/213 (41)
Cough	82/84 (98)	191/213 (90)
Shortness of breath	35/84 (42)	131/210 (62)
Chest pain	38/84 (45)	85/210 (40)
Diarrhoea	11/84 (13)	22/210 (10)
Underlying conditions		
Hypertension	13/84 (15)	36/211 (17)
Cardiac	1/99 (1)	7/222 (3)
Lung disease	0/84 (0)	1/211 (1)
Diabetes	3/84 (4)	24/211 (11)
Cancer	0/99 (0)	3/222 (1)
Tuberculosis	0/84 (0)	17/208 (8)
HIV-infection	13/99 (13)	78/222 (35)
Other **	1/84 (1)	2/211 (1)
SARS-CoV-2 Vaccine		
Pfizer-BioNTech (1 st dose)	12/99 (12)	22/222 (10)
Pfizer-BioNTech (2 nd dose)	12/99 (12)	16/222 (7)
Johnson & Johnson (1 st dose)	11/99 (11)	21/222 (9)
Johnson & Johnson (2 nd dose)	1/99 (1)	1/222 (0)
Unknown	4/99 (4)	2/222 (0)
No vaccine	52/99 (53)	164/222 (74)
Management		
Oxygen therapy	0/83 (0)	110/200 (55)
ICU admission	N/A	0/202 (0)
Ventilation	N/A	3/202 (1)
Outcome***		
Died	0/82 (0)	13/187 (7)

*ILI surveillance not conducted in Gauteng province

**Chronic lung, liver and kidney disease, organ transplant, pregnancy, malnutrition, obesity, tracheostomy, prematurity, seizure, stroke, anaemia, asplenia, burns, Systemic lupus erythematosus, seizures

***Outcome includes patients who are still hospitalised, have been discharged or referred, and those who died

⁵ Pending results: outstanding variant results

⁵⁵ Unable to assign: no lineage assigned due to poor- sequence quality OR low viral load (ct>35) OR variant PCR could not assign variant and no sequencing result

Note: Children may be over-represented amongst hospitalised patients due to the inclusion of a large paediatric hospital in Cape Town.

Of the 13 patients who died, four were in the 20-39 year age group, four were in 40-59 age group and five were ≥60 years; 7/13 (54%) were female.

Methods

SARS-CoV-2 Testing

March 2020 – March 2021: SARS-CoV-2 was detected using the Roche E gene real-time PCR assay (Corman et al. *Euro Surveillance* 2020) with cycle threshold (C_t) <40 interpreted as positive for SARS-CoV-2. From April 2021 to date the laboratory changed to the Allplex™ SARS-CoV-2/FluA/FluB/RSV kit (Seegene Inc., Seoul, South Korea), with positivity assigned if the PCR cycle threshold (C_t) was <40 for ≥1 gene targets (N, S or RdRp).

A confirmed SARS-CoV-2 case is a person of any age enrolled in surveillance with laboratory confirmation of SARS-CoV-2 infection by PCR. Only positive SARS-CoV-2 specimens on PCR are further tested to determine variant/lineage type by variant PCR or genomic sequencing.

Variant PCR

Allplex™ SARS-CoV-2 Variants I PCR detects Alpha and Beta/Gamma variants. The assay was conducted on all SARS-CoV-2-positive samples from 1 March 2020 – 30 June 2021.

Allplex™ SARS-CoV-2 Variants II PCR detects Delta variant and distinguishes Beta from Gamma. The assay was conducted on SARS-CoV-2-positive samples from 1 Jan to 30 June 2021.

Extraction: Total nucleic acids were extracted from 200µl NP/OP samples in universal or viral transport medium using a MagNA Pure 96 automated extractor and DNA/Viral NA Small Volume v2.0 extraction kit (Roche Diagnostics, Mannheim, Germany).

SARS-CoV-2 genomic surveillance

SARS-CoV-2 Whole-Genome Sequencing and Genome Assembly

RNA Extraction

RNA was extracted either manually or automatically in batches, using the QIAamp viral RNA mini kit (QIAGEN, CA, USA) or the Chemagic 360 using the CMG-1049 kit (PerkinElmer, MA, USA). A modification was done on the manual extractions by adding 280 µl per sample, in order to increase yields. 300 µl of each sample was used for automated magnetic bead-based extraction using the Chemagic 360. RNA was eluted in 60 µl of the elution buffer. Isolated RNA was stored at -80 °C prior to use.

PCR and Library Preparation

Sequencing was performed using the Illumina COVIDSeq protocol (Illumina Inc., CA, USA) or nCoV-2019 ARTIC network sequencing protocol v3 (<https://artic.network/ncov-2019>). These are amplicon-based next-generation sequencing approaches. Briefly, for the nCoV-2019 ARTIC network sequencing protocol, the first strand synthesis was carried out on extracted RNA samples using random hexamer primers from the SuperScript IV reverse transcriptase synthesis kit (Life Technologies, CA, USA) or LunaScript RT SuperMix Kit (New England Biolabs (NEB), MA, USA). The synthesized cDNA was amplified using multiplex polymerase chain reactions (PCRs) using ARTIC nCoV-2019 v3 primers. For the COVIDSeq protocol, the first strand synthesis was carried out using random hexamer primers from Illumina and the synthesized cDNA underwent two separate multiplex PCR reactions.

For Illumina sequencing using the nCoV-2019 ARTIC network sequencing protocol, the pooled PCR products underwent bead-based tagmentation using the Nextera Flex DNA library preparation kit (Illumina Inc., CA, USA). The adapter-tagged amplicons were cleaned up using AmpureXP purification beads (Beckman Coulter, High Wycombe, UK) and amplified using one round of PCR. The PCRs were indexed using the Nextera CD indexes (Illumina Inc., CA, USA) according to the manufacturer's instructions. For COVIDSeq sequencing protocol, pooled PCR amplified products were processed for tagmentation and adapter ligation using IDT for Illumina Nextera UD Indexes. Further enrichment and cleanup was performed as per protocols provided by the manufacturer (Illumina Inc., CA, USA). Pooled samples from both COVIDSeq protocol and nCoV-2019 ARTIC network protocol were quantified using Qubit 3.0 or 4.0 fluorometer (Invitrogen Inc., MA, USA) using the Qubit dsDNA High Sensitivity assay according to manufacturer's instructions. The fragment sizes were analyzed using TapeStation 4200 (Invitrogen Inc., MA, USA). The pooled libraries were further normalized to 4nM concentration and 25 µl of each normalized pool containing unique index adapter sets were combined in a new tube. The final library pool was denatured and neutralized with 0.2 N sodium hydroxide and 200 mM Tris-HCL (pH7), respectively. 1.5 pM sample library was spiked with 2% PhiX. Libraries were loaded onto a 300-cycle NextSeq 500/550 HighOutput Kit v2 and run on the Illumina NextSeq 550 instrument (Illumina Inc., CA, USA).

Assembly, Processing and Quality Control of Genomic Sequences

Raw reads from Illumina sequencing were assembled using the Exatype NGS SARS-CoV-2 pipeline v1.6.1, (<https://sars-cov-2.exatype.com/>). The resulting consensus sequence was further manually polished by considering and correcting indels in homopolymer regions that break the open reading frame (probably sequencing errors) using Aliview v1.27, (<http://orombunkar.se/aliview/>) (Larsson, 2014). Mutations resulting in mid-gene stop codons and frameshifts were reverted to wild type. All assemblies determined to have acceptable quality (defined as having at least 1 000 000 reads and at least 40 % 10 X coverage) were deposited on GISAID (<https://www.gisaid.org/>) (Elbe & Buckland-Merrett, 2017; Shu & McCauley, 2017).

Classification of Lineage, Clade and Associated Mutations

Assembled genomes were assigned lineages using the 'Phylogenetic Assignment of Named Global Outbreak Lineages' (PANGOLIN) software suite (<https://github.com/hCoV-2019/pangolin>) (Rambaut et al., 2020), a tool used for dynamic SARS-CoV-2 lineage classification. The SARS-CoV-2 genomes in our dataset were also classified using the clade classification proposed by NextStrain (<https://nextstrain.org/>), a tool built for real-time tracking of the pathogen evolution (Hadfield et al., 2018).